

Collapsing behaviour of a singular diffusion equation

Kin Ming Hui

Institute of Mathematics, Academia Sinica,
Nankang, Taipei, 11529, Taiwan, R. O. C.

May 27, 2011

Abstract

Let $0 \leq u_0(x) \in L^1(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ be such that $u_0(x) = u_0(|x|)$ for all $|x| \geq r_1$ and is monotone decreasing for all $|x| \geq r_1$ for some constant $r_1 > 0$ and $\text{ess inf}_{\overline{B_{r_1}}(0)} u_0 \geq \text{ess sup}_{\mathbb{R}^2 \setminus B_{r_2}(0)} u_0$ for some constant $r_2 > r_1$. Then under some mild decay conditions at infinity on the initial value u_0 we will extend the result of P. Daskalopoulos, M.A. del Pino and N. Sesum [DP2], [DS2], and prove the collapsing behaviour of the maximal solution of the equation $u_t = \Delta \log u$ in $\mathbb{R}^2 \times (0, T)$, $u(x, 0) = u_0(x)$ in $\mathbb{R}^2$, near its extinction time $T = \int_{\mathbb{R}^2} u_0 dx / 4\pi$ without using the Hamilton-Yau Harnack inequality.

Key words: collapsing behaviour, maximal solution, singular diffusion equation
AMS Mathematics Subject Classification: Primary 35B40 Secondary 35K57, 35K65

0 Introduction

Recently there is a lot of study on the equation,

$$\begin{cases} u_t = \Delta \log u, u > 0, & \text{in } \mathbb{R}^2 \times (0, T) \\ u(x, 0) = u_0 & \text{in } \mathbb{R}^2 \end{cases} \quad (0.1)$$

by P. Daskalopoulos, M.A. del Pino and N. Sesum [DP1], [DP2], [DS1], [DS2], S.Y. Hsu [Hs1], [Hs2], [Hs3], [Hs4], J.R. Esteban, A. Rodriguez and J.L. Vazquez [ERV1], [ERV2], K.M. Hui [Hu1], [Hu2], and L.F. Wu [W1], [W2], etc. Equation (0.1) arises in the model of thin films of fluid spreading on a solid surface [G] when the fourth order term is neglected. It also arises in the study of the Ricci flow on surfaces [W1], [W2], and as the singular limit of the porous medium equation,

$$\begin{cases} u_t = \Delta \left(\frac{u^m}{m} \right), u > 0, & \text{in } \mathbb{R}^2 \times (0, T) \\ u(x, 0) = u_0(x) & \text{in } \mathbb{R}^2 \end{cases}$$

as $m \rightarrow 0$ [Hu2], [ERV2]. In [DP1] and [Hu1] P. Daskalopoulos and M.A. del Pino and K.M. Hui proved that corresponding to each $0 \leq u_0 \in L^p(\mathbb{R}^2) \cap L^1(\mathbb{R}^2)$, $p > 1$, and $2 \leq f \in L^1(0, \infty)$ there exists a classical solution u of (0.1) in $\mathbb{R}^n \times (0, T)$ satisfying the mass loss equation,

$$\int_{\mathbb{R}^2} u(x, t) dx = \int_{\mathbb{R}^2} u_0 dx - 2\pi \int_0^t f(s) ds \quad \forall 0 \leq t < T \quad (0.2)$$

where $T > 0$ given by

$$\int_{\mathbb{R}^2} u_0 dx = 2\pi \int_0^T f(s) ds \quad (0.3)$$

is the extinction time for the solution u . Note that the maximal solution of (0.1) is the solution of (0.1) that satisfies (0.2) with $f \equiv 2$. When $f \equiv \gamma > 2$ is a constant, the asymptotic behaviour of the solution u is studied by S.Y. Hsu in [Hs3] and [Hs4]. In [Hs3] S.Y. Hsu proved that if u_0 is radially symmetric and monotone decreasing and u is the solution of (0.1) in $\mathbb{R}^2 \times (0, T)$ that satisfies (0.2), (0.3), with $f \equiv 4$, then the rescaled solution

$$v(x, s) = \frac{u(x, t)}{T - t}, \quad s = -\log(T - t)$$

will converge uniformly on every compact subsets of $\mathbb{R}^2$ to the function

$$\frac{8\lambda}{(\lambda + |x|^2)^2}$$

as $s \rightarrow \infty$ for some constant $\lambda > 0$. For the general case $f \equiv \gamma > 2$ S.Y. Hsu [Hs4] proved that if u_0 is radially symmetric and monotone decreasing and u is the solution of (0.1) in $\mathbb{R}^2 \times (0, T)$ that satisfies (0.2) and (0.3) with $f \equiv \gamma$ and

$$\lim_{r \rightarrow \infty} \frac{ru_r(r, t)}{u(r, t)} = -\gamma \quad \text{uniformly on } [a, b] \quad \forall 0 < a < b < T$$

where T is given by (0.3) with $f \equiv \gamma$, then there exist unique constants $\alpha > 0$, $\beta > -1/2$, $\alpha = 2\beta + 1$, depending only on γ such that the rescaled function

$$v(y, s) = \frac{u(y/(T - t)^\beta, t)}{(T - t)^\alpha}$$

where

$$s = -\log(T - t)$$

will converge uniformly on every compact subset of $\mathbb{R}^2$ to $\phi_{\lambda, \beta}(y)$ for some constant $\lambda > 0$ as $s \rightarrow \infty$ where $\phi_{\lambda, \beta}(y) = \phi_{\lambda, \beta}(|y|)$ is radially symmetric and satisfies the following O.D.E.

$$\frac{1}{r} \left(\frac{r\phi'}{\phi} \right)' + \alpha\phi + \beta r\phi' = 0 \quad \text{in } (0, \infty)$$

with

$$\phi(0) = 1/\lambda, \phi'(0) = 0.$$

However as observed by J.R. King [K] using formal asymptotic analysis as t approaches T the vanishing behaviour for the finite mass solution of (0.1) which corresponds to the case $f \equiv 2$ is very different from the vanishing behaviour for the case $f \equiv \gamma > 2$. J.R. King find that for compactly supportly finite mass initial value the maximal solution of (0.1) behaves like

$$\frac{(T-t)^2}{\frac{T}{2}|x|^2 + e^{\frac{2T}{T-t}}}$$

in the inner region $(T-t) \log |x| \leq T$ and behaves like

$$\frac{2t}{|x|^2(\log |x|)^2}$$

in the outer region $(T-t) \log |x| \geq T$ as $t \nearrow T$. In [DP2] P. Daskalopoulos and M.A. del Pino give a rigorous proof of an extension of this formal result for radially symmetric initial value $u_0(r)$ satisfying the conditions,

$$u_0(x) = u_0(|x|) \text{ is decreasing on } r = |x| \geq r_1 \text{ for some constant } r_1 > 0 \quad (0.4)$$

$$u_0(x) = \frac{2\mu}{|x|^2(\log |x|)^2}(1 + o(1)) \quad \text{as } |x| \rightarrow \infty, \quad (0.5)$$

for some constant $\mu > 0$ and

$$R_0(x) := -\frac{\Delta \log u_0}{u_0} \geq -\frac{1}{\mu} \quad \text{on } \mathbb{R}^2. \quad (0.6)$$

Note that (0.1) is equivalent to the Ricci flow equation ([W2])

$$\frac{\partial}{\partial t} g_{ij} = -2R_{ij} \quad \text{in } \mathbb{R}^2 \times (0, T)$$

where $g_{ij}(t) = g_{ij}(x, t) = u(x, t)\delta_{ij}$ is an evolving metric on $\mathbb{R}^2$ and $R_{ij}(x, t)$ is the Ricci curvature of the metric $g_{ij}(t)$. Then the scalar curvature $R(x, t)$ with respect to the metric $g_{ij}(x, t)$ is given by

$$R(x, t) = -\frac{\Delta \log u}{u}.$$

Thus (0.6) says that the initial scalar curvature is greater than or equal to $-1/\mu$ on $\mathbb{R}^2$.

In [DS2] P. Daskalopoulos and N. Sesum extend the result to the case of compactly supported $0 \leq u_0 \in L^1(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$. However their proof of the behaviour of the

maximal solution in the outer region near the extinction time is very difficult and uses the Hamilton-Yau Harnack inequality [HY].

In this paper we will extend their result to the case of initial value $0 \leq u_0 \in L^1(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ that satisfies (0.4), (0.5), and (0.6) for some constant $\mu \geq 0$ with the right hand side being replaced by $-\infty$ if $\mu = 0$ and

$$\operatorname{ess\,inf}_{\overline{B}_{r_1}(0)} u_0 \geq \operatorname{ess\,sup}_{\mathbb{R}^2 \setminus B_{r_2}(0)} u_0 \quad (0.7)$$

for some constant $r_2 > r_1$. Note that (0.7) is automatically satisfied if u_0 has compact support in $\mathbb{R}^2$. We will prove the behaviour of the maximal solution in the outer region near the extinction time by elementary method without using the difficult Hamilton-Yau Harnack inequality for surfaces [HY]. We will also prove the behaviour of the maximal solution in the inner region as the extinction time is approached.

In this paper we will assume that the initial value u_0 satisfies $0 \leq u_0 \in L^1(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$, (0.4), (0.5), (0.6) and (0.7) for some constants $r_2 > r_1 > 0$ and $\mu \geq 0$ with the right hand side being replaced by $-\infty$ if $\mu = 0$. We will assume that u is the maximal solution of (0.1) in $\mathbb{R}^2 \times (0, T)$ that satisfies (0.2) with $f \equiv 2$ and

$$T = \frac{1}{4\pi} \int_{\mathbb{R}^2} u_0 \, dx$$

for the rest of the paper. For any $x_0 \in \mathbb{R}^2$ and $r_0 > 0$ let $B_{r_0}(x_0) = \{x \in \mathbb{R}^2 : |x - x_0| < r_0\}$ and $B_{r_0} = B_{r_0}(0)$. Note that by [Hu1] u satisfies the Aronson-Bénilan inequality,

$$\frac{u_t}{u} \leq \frac{1}{t} \quad \text{in } \mathbb{R}^2 \times (0, T). \quad (0.8)$$

The plan of the paper is as follows. In section 1 we will prove the behaviour of the maximal solution in the inner region. In section two we will prove the behaviour of the maximal solution in the outer region.

1 Inner region behaviour

In this section we will prove the behaviour of the maximal solution in the inner region near the extinction time using a modification of the argument of [DP2] and [DS2].

Lemma 1.1. *The solution u satisfies*

$$u(x, t) \geq u(y, t)$$

for any $t \in (0, T)$ and $x, y \in \mathbb{R}^2$ such that $|y| \geq |x| + 2r_2$.

Proof: We will use a modification of the proof of Lemma 2.1 of [CF] to prove the lemma.

For any $x^0, y^0 \in \mathbb{R}^2$ such that $|y^0| \geq |x^0| + 2r_2$ let Π be the hyperplane of points in $\mathbb{R}^2$ which are equidistance from x^0 and y^0 . Then (cf. Lemma 2.1 of [CF]),

$$\Pi = \{x \in \mathbb{R}^2 : x \cdot (x^0 - y^0) = \frac{1}{2}(x^0 + y^0) \cdot (x^0 - y^0)\}$$

and

$$\text{dist}(\Pi, \{0\}) = \frac{1}{2} \cdot \frac{|y^0|^2 - |x^0|^2}{|x^0 - y^0|} \geq \frac{1}{2} \cdot \frac{|y^0|^2 - |x^0|^2}{|x^0| + |y^0|} \geq \frac{1}{2}(|y^0| - |x^0|) \geq r_2. \quad (1.1)$$

We write $\mathbb{R}^2 \setminus \Pi = \Pi_+ \cup \Pi_-$ where Π_+ and Π_- are the two half-spaces with respect to Π with $0 \in \Pi_-$. By (1.1) $\{x^0\} \cup B_{r_2} \subset \Pi_-$ and $y^0 \in \Pi_+$. By rotation we may assume without loss of generality that

$$\begin{cases} \Pi = \{(x_1, x_2) \in \mathbb{R}^2 : x_1 = a_0\} \\ \Pi_- = \{(x_1, x_2) \in \mathbb{R}^2 : x_1 < a_0\} \\ \Pi_+ = \{(x_1, x_2) \in \mathbb{R}^2 : x_1 > a_0\} \end{cases}$$

where $a_0 = \text{dist}(\Pi, \{0\})$. For any $x = (x_1, x_2) \in \Pi_-$ let $\tilde{x} = (2a_0 - x_1, x_2)$ be the reflection point of x about Π . Then if $x \in \overline{B}_{r_1}$, by (0.7) and (1.2)

$$u_0(x) \geq u_0(\tilde{x}). \quad (1.2)$$

If $x \in \Pi_- \setminus \overline{B}_{r_1}$, then

$$\begin{aligned} 2a_0 - x_1 > |x_1| &\Rightarrow |\tilde{x}| \geq |x| > r_1 \\ &\Rightarrow u_0(x) \geq u_0(\tilde{x}) \quad (\text{by (0.4)}). \end{aligned}$$

Hence (1.2) holds for any $x \in \Pi_-$. By the maximum principle for the equation

$$u_t = \Delta \log u$$

in the half-space Π_- (cf. Lemma 2.5 of [ERV2]),

$$u(x, t) \geq u(\tilde{x}, t) \quad \forall x \in \Pi_-, 0 < t < T.$$

Hence

$$u(x^0, t) \geq u(y^0, t)$$

and the lemma follows. □

By Lemma 1.1 for any $0 < t < T$ there exists $x_t \in \overline{B}_{2r_2}$ such that

$$u(x_t, t) = \max_{x \in \mathbb{R}^2} u(x, t). \quad (1.3)$$

Similar to [DP2] we let

$$\overline{u}(x, \tau) = \tau^2 u(x, t), \quad \tau = \frac{1}{T - t}, \tau > 1/T. \quad (1.4)$$

Then $\bar{u}$ satisfies [DP2],

$$\bar{u}_\tau = \Delta \log \bar{u} + \frac{2\bar{u}}{\tau} \quad \text{in } \mathbb{R}^2 \times (1/T, \infty). \quad (1.5)$$

Let $R_{\max}(t) = \max_{x \in \mathbb{R}^2} R(x, t)$ and let $W(t)$ be the width function with respect to the metric $g_{ij}(t)$ as defined by P. Daskalopoulos and R. Hamilton in [DH]. We recall a result of [DH].

Theorem 1.2. [DH] *There exist positive constants $c > 0$ and $C > 0$ such that*

$$(i) \quad c(T - t) \leq W(t) \leq C(T - t)$$

$$(ii) \quad \frac{c}{(T-t)^2} \leq R_{\max}(t) \leq \frac{C}{(T-t)^2}$$

hold for any $0 < t < T$.

Let

$$\bar{R}(x, \tau) = -\frac{\Delta \log \bar{u}}{\bar{u}}.$$

Similar to the argument on P.862–863 of [DP2] by Theorem 1.2, (1.4) and the Aronson-Bénlian inequality (0.8),

$$\begin{aligned} -\frac{1}{\tau^2 t} &\leq \bar{R}(x, \tau) \leq C \quad \forall (x, \tau) \in \mathbb{R}^2 \times (1/T, \infty), t = T - (1/\tau) \\ \Rightarrow \quad -\frac{2}{\tau^2 T} &\leq \bar{R}(x, \tau) \leq C \quad \forall (x, \tau) \in \mathbb{R}^2 \times (2/T, \infty). \end{aligned} \quad (1.6)$$

By (1.5) and (1.6),

$$C \geq \bar{R}(x, \tau) = -\frac{\Delta \log \bar{u}}{\bar{u}} = \frac{-\bar{u}_\tau + \frac{2\bar{u}}{\tau}}{\bar{u}} = -\frac{\bar{u}_\tau}{\bar{u}} + \frac{2}{\tau} \geq -\frac{2}{\tau^2 T} \quad \text{in } \mathbb{R}^2 \times (2/T, \infty) \quad (1.7)$$

$$\Rightarrow \quad \frac{2}{\tau} + \frac{2}{\tau^2 T} \geq \frac{\bar{u}_\tau}{\bar{u}} \geq -C + \frac{2}{\tau} \geq -C \quad \text{in } \mathbb{R}^2 \times (2/T, \infty). \quad (1.8)$$

Theorem 1.3. *For any sequence $\{\tau_k\}_{k=1}^\infty$, $\tau_k \rightarrow \infty$ as $k \rightarrow \infty$, let $x_{t_k} \in \bar{B}_{2r_2}$ be given by (1.3) with $t = t_k$ and*

$$\bar{u}_k(y, \tau) = \alpha_k \bar{u}(\alpha_k^{\frac{1}{2}} y + x_{t_k}, \tau + \tau_k), \quad y \in \mathbb{R}^2, \tau > -\tau_k + T^{-1} \quad (1.9)$$

where

$$t_k = T - \tau_k^{-1} \quad \forall k \in \mathbb{Z}^+ \quad (1.10)$$

and

$$\alpha_k = 1/\bar{u}(x_{t_k}, \tau_k). \quad (1.11)$$

Then $\{\bar{u}_k\}_{k=1}^\infty$ has a subsequence $\{\bar{u}_{k_i}\}_{i=1}^\infty$ that converges uniformly on $C^\infty(K)$ for any compact set $K \subset \mathbb{R}^2 \times (-\infty, \infty)$ as $i \rightarrow \infty$ to a positive solution U of equation

$$U_\tau = \Delta \log U \quad \text{in } \mathbb{R}^2 \times (-\infty, \infty) \quad (1.12)$$

with uniformly bounded non-negative scalar curvature and uniformly bounded width on $\mathbb{R}^2 \times (-\infty, \infty)$ with respect to the metric $\tilde{g}_{ij}(t) = U(\cdot, t)\delta_{ij}$.

Proof: We first observe that by (1.5),

$$\bar{u}_{k,\tau} = \Delta \log \bar{u}_k + \frac{2\bar{u}_k}{\tau + \tau_k} \quad \text{in } \mathbb{R}^2 \times (-\tau_k + (1/T), \infty) \quad (1.13)$$

with

$$\bar{u}_k(0, 0) = 1 \quad \text{and} \quad \bar{u}_k(y, 0) \leq 1 \quad \forall y \in \mathbb{R}^2. \quad (1.14)$$

Since $(\log \bar{u}_k)_\tau = (\log \bar{u})_\tau$, by (1.8),

$$-C \leq \frac{\bar{u}_{k,\tau}}{\bar{u}_k} \leq \frac{2}{\tau + \tau_k} + \frac{2}{(\tau + \tau_k)^2 T} \leq \frac{3T}{2} \quad \text{in } \mathbb{R}^2 \times (2/T - \tau_k, \infty). \quad (1.15)$$

For any $-\infty < a < 0 < b < \infty$ we choose $k_0 \in \mathbb{Z}^+$ such that $-\tau_k + (2/T) < a$ for any $k \geq k_0$. Then by (1.14) and (1.15) there exists a constant $M_1 > 0$ such that

$$\bar{u}_k(x, \tau) \leq M_1 \quad x \in \mathbb{R}^2, a \leq \tau \leq b, k \geq k_0. \quad (1.16)$$

By (1.15) and (1.16) there exists a constant $C > 0$ such that

$$|\bar{u}_{k,\tau}(x, \tau)| \leq CM_1 \quad x \in \mathbb{R}^2, a \leq \tau \leq b, k \geq k_0. \quad (1.17)$$

To complete the proof of the theorem we need the following two technical lemmas.

Lemma 1.4. *There exists a constant $C_1 > 0$ such that*

$$\begin{aligned} -C_1 R_1^2 M_1 + \log \left(\frac{1}{\bar{u}_k(x_0, \tau)} \right) &\leq \frac{1}{|B_{R_1}|} \int_{B_{R_1}(x_0)} \log \left(\frac{1}{\bar{u}_k(x, \tau)} \right) dx \\ &\leq C_1 R_1^2 M_1 + \log \left(\frac{1}{\bar{u}_k(x_0, \tau)} \right) \end{aligned} \quad (1.18)$$

holds for any $R_1 > 0$, $x_0 \in \mathbb{R}^2$, $a \leq \tau \leq b$ and $k \geq k_0$.

Proof: We will use a modification of the proof of Lemma 6 of [V] and Lemma 2.6 of [Hu1] to proof the lemma. Let

$$G_{R_1}(x) = \log(R_1/|x - x_0|) + \frac{1}{2}R_1^{-2}(|x - x_0|^2 - R_1^2)$$

be the Green function for $B_{R_1}(x_0)$. Then $G_{R_1} \geq 0$ and $\Delta G_{R_1} = 2R_1^{-2} - 2\pi\delta_0$ where δ_0 is the delta mass at the origin. By (1.13),

$$\begin{aligned}
& \int_{B_{R_1}(x_0)} G_{R_1}(x) \bar{u}_{k,\tau}(x, \tau) dx \\
&= \int_{B_{R_1}(x_0)} G_{R_1}(x) \left(\Delta \log \bar{u}_k(x, \tau) + \frac{2\bar{u}_k(x, \tau)}{\tau + \tau_k} \right) dx \\
&= 2\pi \left(\log \left(\frac{1}{\bar{u}_k(x_0, \tau)} \right) - \frac{1}{|B_{R_1}|} \int_{B_{R_1}(x_0)} \log \left(\frac{1}{\bar{u}_k(x, \tau)} \right) dx \right) \\
&\quad + \frac{2}{\tau + \tau_k} \int_{B_{R_1}(x_0)} G_{R_1}(x) \bar{u}_k(x, \tau) dx \quad \forall \tau \geq -\tau_k + \frac{2}{T}, k \geq k_0. \tag{1.19}
\end{aligned}$$

Since

$$\int_{B_{R_1}(x_0)} G_{R_1}(x) dx \leq CR_1^2,$$

by (1.16), (1.17), and (1.19) we get (1.18) and the lemma follows. $\square$

Lemma 1.5. *For any $R_1 > 0$ there exists a constant $C_2 > 0$ such that*

$$\sup_{\substack{|y| \leq R_1 \\ a \leq \tau_1 \leq b}} \bar{u}_k(y, \tau_1)^9 \leq C_2 \inf_{\substack{|x| \leq R_1 \\ a \leq \tau_2 \leq b}} \bar{u}_k(x, \tau_2) \quad \forall k \geq k_0.$$

Proof: Let $|x_0|, |y_0| \leq R_1$, $\tau_1, \tau_2 \in [a, b]$ and $k \geq k_0$. Since $B_{R_1}(x_0) \subset B_{3R_1}(y_0)$, by Lemma 1.4,

$$\begin{aligned}
\log \left(\frac{1}{\bar{u}_k(x_0, \tau_1)} \right) &\leq \frac{1}{|B_{R_1}|} \int_{B_{R_1}(x_0)} \log \left(\frac{1}{\bar{u}_k(x, \tau_1)} \right) dx + C_1 M_1 R_1^2 \\
&\leq \frac{9}{|B_{3R_1}|} \int_{B_{3R_1}(y_0)} \log \left(\frac{1}{\bar{u}_k(x, \tau_1)} \right) dx + C_1 M_1 R_1^2 \\
&\leq 9 \log \left(\frac{1}{\bar{u}_k(y_0, \tau_1)} \right) + C' M_1 R_1^2
\end{aligned}$$

for some constants $C_1 > 0$ and $C' = 82C_1$. Hence

$$\frac{1}{\bar{u}_k(x_0, \tau_1)} \leq \frac{e^{C' M_1 R_1^2}}{\bar{u}_k(y_0, \tau_1)^9} \Rightarrow \bar{u}_k(y_0, \tau_1)^9 \leq e^{C' M_1 R_1^2} \bar{u}_k(x_0, \tau_1). \tag{1.20}$$

Now by (1.15) there exists a constant $C > 0$ such that

$$\bar{u}_k(x, \tau'_1) \leq C \bar{u}_k(x, \tau'_2) \quad \forall x \in \mathbb{R}^2, \tau'_1, \tau'_2 \in [a, b], k \geq k_0. \tag{1.21}$$

By (1.20) and (1.21),

$$\bar{u}_k(y_0, \tau_1)^9 \leq C_2 \bar{u}_k(x_0, \tau_2)$$

holds for some constant $C_2 > 0$ and the lemma follows. $\square$

We will now continue with the proof of Theorem 1.3. By (1.14) and Lemma 1.5 for any $R_1 > 0$ there exist constants $C_3 > 0$ and $C_4 > 0$ such that

$$C_3 \leq \bar{u}_k(x, \tau_1) \leq C_4 \quad \forall |x| \leq R_0, a \leq \tau \leq b, k \geq k_0.$$

Hence the equation (1.13) for $\bar{u}_k$ is uniformly parabolic on $\bar{B}_{R_1} \times [a, b]$ for all $k \geq k_0$. By the parabolic Schauder estimates [LSU] $\bar{u}_k$ are uniformly Hölder continuous in $C^{2\gamma, 1\gamma}(\bar{B}_{R_1} \times [a, b])$ for any $\gamma \in \mathbb{Z}^+$. By the Ascoli theorem and a diagonalization argument the sequence $\{\bar{u}_k\}_{k=1}^\infty$ has a subsequence which we may assume without loss of generality to be the sequence itself that converges uniformly in $C^\infty(K)$ as $k \rightarrow \infty$ for any compact set $K \subset \mathbb{R}^2 \times (-\infty, \infty)$ to some positive function U that satisfies (1.12).

Let

$$\bar{R}_k = -\frac{\Delta \log \bar{u}_k}{\bar{u}_k}. \quad (1.22)$$

Then $\bar{R}_k$ converges uniformly on every compact subset of $\mathbb{R}^2 \times (-\infty, \infty)$ as $k \rightarrow \infty$ to the scalar curvature $\tilde{R} = -(\Delta \log U)/U$ of the metric $\tilde{g}_{ij}(\tau) = U(\cdot, \tau)\delta_{ij}$. Note that by (1.6),

$$\begin{aligned} -\frac{2}{(\tau + \tau_k)^2 T} &\leq \bar{R}_k(y, \tau) \leq C \quad \forall (y, \tau) \in \mathbb{R}^2 \times (2/T - \tau_k, \infty) \\ \Rightarrow 0 &\leq \tilde{R}(y, \tau) \leq C \quad \forall (y, \tau) \in \mathbb{R}^2 \times (-\infty, \infty) \quad \text{as } k \rightarrow \infty. \end{aligned}$$

Finally similar to the argument on P.10 of [DS2] by Theorem 1.2 and an approximation argument the width function with respect to the metric $\tilde{g}_{ij}(\tau) = U(\cdot, \tau)\delta_{ij}$ is uniformly bounded on $\mathbb{R}^2 \times (-\infty, \infty)$. $\square$

Theorem 1.6. *For any sequence $\{\tau_k\}_{k=1}^\infty$, $\tau_k \rightarrow \infty$ as $k \rightarrow \infty$, let $\bar{u}_k$ be given by (1.9). Then $\{\bar{u}_k\}_{k=1}^\infty$ has a subsequence $\{\bar{u}_{k_i}\}_{i=1}^\infty$ that converges uniformly in $C^\infty(K)$ for any compact set $K \subset \mathbb{R}^2 \times (-\infty, \infty)$ as $i \rightarrow \infty$ to*

$$U(y, \tau) = \frac{1}{\lambda|y|^2 + e^{4\lambda\tau}}. \quad (1.23)$$

for some constant $\lambda > 0$.

Proof: By Theorem 1.3 $\{\bar{u}_k\}_{k=1}^\infty$ has a subsequence $\{\bar{u}_{k_i}\}_{i=1}^\infty$ that converges uniformly in $C^\infty(K)$ for any compact set $K \subset \mathbb{R}^2 \times (-\infty, \infty)$ as $i \rightarrow \infty$ to a solution $U(y, \tau)$ of (1.12). By Theorem 1.3 and the result of [DS1],

$$U(y, \tau) = \frac{2}{\beta(|y - y_0|^2 + \delta e^{2\beta\tau})} \quad (1.24)$$

for some $y_0 \in \mathbb{R}^2$ and constants $\beta > 0$, $\delta > 0$. Since $\bar{u}_{k_i}$ converges uniformly on every compact subset of $\mathbb{R}^2 \times (-\infty, \infty)$ to $U(y, \tau)$ and $\bar{u}_k(y, 0)$ attains its maximum at $y = 0$, $U(y, 0)$ will attain its maximum at $y = 0$. Hence $y_0 = 0$. By (1.14),

$$U(0, 0) = 1 \quad \Rightarrow \quad 1 = \frac{2}{\beta\delta}. \quad (1.25)$$

By (1.24) and (1.25) we get (1.23) with $\lambda = \beta/2 > 0$. $\square$

We now let

$$\alpha(\tau) = 1/\bar{u}(x_t, \tau) \quad (1.26)$$

where $\tau = 1/(T - t)$, $\tau > 1/T$, and $x_t \in \bar{B}_{2r_2}$ satisfies (1.3).

Lemma 1.7. *There exist constants $\delta > 0$ and $\tau_0 > 1/T$ such that*

$$\liminf_{\delta' \rightarrow 0^+} \left(\frac{\log \alpha(\tau) - \log \alpha(\tau - \delta')}{\delta'} \right) > \delta \quad \forall \tau \geq \tau_0. \quad (1.27)$$

Proof: Suppose (1.27) does not hold. Then there exist a sequence of positive numbers $\{\delta_k\}_{k=1}^\infty$, $\delta_k \rightarrow 0$ as $k \rightarrow \infty$, and a sequence $\{\tau_k\}_{k=1}^\infty$, $\tau_k > 1/T$ for all $k \in \mathbb{Z}^+$ and $\tau_k \rightarrow \infty$ as $k \rightarrow \infty$, such that

$$\liminf_{\delta' \rightarrow 0^+} \left(\frac{\log \alpha(\tau_k) - \log \alpha(\tau_k - \delta')}{\delta'} \right) \leq \delta_k \quad \forall k \in \mathbb{Z}^+.$$

Hence for each $k \in \mathbb{Z}^+$ there exists a sequence of positive numbers $\{\delta_{k,j}\}_{j=1}^\infty$, $\delta_{k,j} \rightarrow 0$ as $j \rightarrow \infty$, such that

$$\frac{\log \alpha(\tau_k) - \log \alpha(\tau_k - \delta_{k,j})}{\delta_{k,j}} < 2\delta_k \quad \forall k, j \in \mathbb{Z}^+. \quad (1.28)$$

Let t_k be given by (1.10) and let $t_{k,j} = T - (\tau_k - \delta_{k,j})^{-1}$. Let x_{t_k} and $x_{t_{k,j}}$ be given by (1.3) with $t = t_k, t_{k,j}$ respectively. Then by (1.26) and (1.28),

$$\frac{\log \bar{u}(x_{t_{k,j}}, \tau_k - \delta_{k,j}) - \log \bar{u}(x_{t_k}, \tau_k)}{\delta_{k,j}} < 2\delta_k \quad \forall k, j \in \mathbb{Z}^+. \quad (1.29)$$

Since

$$\bar{u}(x_{t_k}, \tau_k - \delta_{k,j}) \leq \max_{z \in \mathbb{R}^2} \bar{u}(z, \tau_k - \delta_{k,j}) = \bar{u}(x_{t_{k,j}}, \tau_k - \delta_{k,j}),$$

by (1.29),

$$\begin{aligned} & \frac{\log \bar{u}(x_{t_k}, \tau_k - \delta_{k,j}) - \log \bar{u}(x_{t_k}, \tau_k)}{\delta_{k,j}} < 2\delta_k \quad \forall k, j \in \mathbb{Z}^+ \\ \Rightarrow & \frac{\bar{u}_\tau}{\bar{u}}(x_{t_k}, \tau_k) \geq -2\delta_k \quad \forall k \in \mathbb{Z}^+ \quad \text{as } j \rightarrow \infty \\ \Rightarrow & \frac{\Delta \log \bar{u}}{\bar{u}}(x_{t_k}, \tau_k) + \frac{2}{\tau_k} \geq -2\delta_k \quad \forall k \in \mathbb{Z}^+ \quad (\text{by (1.5)}) \\ \Rightarrow & -\bar{R}(x_{t_k}, \tau_k) + \frac{2}{\tau_k} \geq -2\delta_k \quad \forall k \in \mathbb{Z}^+. \end{aligned} \quad (1.30)$$

Let $\bar{u}_k$ be given by (1.9) with $\alpha_k = \alpha(\tau_k)$ and $\bar{R}_k$ be given by (1.22). Since $\bar{R}_k(0, 0) = \bar{R}(x_{t_k}, \tau_k)$, by (1.30),

$$\bar{R}_k(0, 0) \leq 2\delta_k + \frac{2}{\tau_k} \quad \forall k \in \mathbb{Z}^+. \quad (1.31)$$

By Theorem 1.3 and Theorem 1.6 $\bar{u}_k$ has a subsequence which we may assume without loss of generality to be the sequence itself that converges uniformly on $C^\infty(K)$ for any compact set $K \subset \mathbb{R}^2 \times (-\infty, \infty)$ as $k \rightarrow \infty$ to some function $U(y, \tau)$ given by (1.23) for some $\lambda > 0$. Then

$$\bar{R}_k(0, 0) \rightarrow -\frac{\Delta \log U}{U}(0, 0) = 4\lambda \quad \text{as } k \rightarrow \infty. \quad (1.32)$$

Letting $k \rightarrow \infty$ in (1.31), by (1.32) we get

$$4\lambda \leq 0.$$

Since $\lambda > 0$, contradiction arises. Hence there exist constants $\delta > 0$ and $\tau_0 > 1/T$ such that (1.27) holds. $\square$

Corollary 1.8. *Let $\delta > 0$ and $\tau_0 > 1/T$ be as given by Lemma 1.7. Then*

$$\alpha(\tau) \geq \alpha(\tau_0) e^{\delta(\tau - \tau_0)} \quad \forall \tau \geq \tau_0.$$

Proof: By Lemma 1.7 there exists constants $\delta > 0$ and $\tau_0 > 1/T$ such that (1.27) holds. Let $\tau > \tau_0$. By (1.27) there exists a constant $\delta'_0 > 0$ such that

$$\log \alpha(\tau) - \log \alpha(\tau - \delta') > \delta \delta' \quad \forall 0 < \delta' \leq \delta'_0. \quad (1.33)$$

Let

$$\delta_0 = \sup\{\delta_1 > 0 : \log \alpha(\tau) - \log \alpha(\tau - \delta') \geq \delta \delta' \quad \forall 0 < \delta' \leq \delta_1\}.$$

Then by (1.33) $\delta_0 \geq \delta'_0$. We claim that $\delta_0 \geq \tau - \tau_0$. Suppose not. Then $\delta_0 < \tau - \tau_0$. By continuity,

$$\log \alpha(\tau) - \log \alpha(\tau - \delta') \geq \delta \delta' \quad \forall 0 < \delta' \leq \delta_0. \quad (1.34)$$

Since $\tau - \delta_0 > \tau_0$, by (1.27) there exists a constant $\delta'_1 > 0$ such that

$$\log \alpha(\tau - \delta_0) - \log \alpha(\tau - \delta_0 - \delta') > \delta \delta' \quad \forall 0 < \delta' \leq \delta'_1. \quad (1.35)$$

By (1.34) and (1.35),

$$\begin{aligned} & \log \alpha(\tau) - \log \alpha(\tau - (\delta_0 + \delta')) > \delta(\delta_0 + \delta') \quad \forall 0 < \delta' \leq \delta'_1 \\ \Rightarrow & \log \alpha(\tau) - \log \alpha(\tau - \delta') \geq \delta \delta' \quad \forall 0 < \delta' \leq \delta_0 + \delta'_1. \end{aligned}$$

This contradicts the definition of δ_0 . Hence $\delta_0 \geq \tau - \tau_0$. Thus

$$\begin{aligned} & \log \alpha(\tau) \geq \log \alpha(\tau_0) + \delta(\tau - \tau_0) \quad \forall \tau \geq \tau_0 \\ \Rightarrow & \alpha(\tau) \geq \alpha(\tau_0) e^{\delta(\tau - \tau_0)} \quad \forall \tau \geq \tau_0 \end{aligned}$$

and the corollary follows. $\square$

Corollary 1.9. *Let α_k be as in Theorem 1.3. Then $\alpha_k \rightarrow \infty$ as $k \rightarrow \infty$.*

Lemma 1.10. *Let τ_k , τ_{k_i} , α_k , $U(y, \tau)$ and $\lambda > 0$ be as in Theorem 1.3 and Theorem 1.6. Let*

$$q_k(y, \tau) = \alpha_k \bar{u}(\alpha_k^{-\frac{1}{2}} y, \tau + \tau_k) \quad (1.36)$$

where $\bar{u}$ is given by (1.4). Then $q_{k_i}(y, \tau)$ converges uniformly in $C^\infty(K)$ for every compact set $K \subset \mathbb{R}^2$ to the function $U(y, \tau)$ as $\tau \rightarrow \infty$.

Proof: Let $\bar{u}_k(y, \tau)$ be given by (1.9) with x_{t_k} , t_k , given by (1.3) and (1.10). Then by Theorem 1.3 and Theorem 1.6 $\bar{u}_k(y, \tau)$ has a subsequence which we may assume without loss of generality to be the sequence itself that converges uniformly on $C^\infty(K)$ for every compact set $K \subset \mathbb{R}^2$ to the function $U(y, \tau)$ given by (1.23) for some $\lambda > 0$ as $k \rightarrow \infty$.

By Corollary 1.9 there exists $k_0 \in \mathbb{Z}^+$ such that $\alpha_k \geq 1$ for all $k \geq k_0$. Let $K \subset \mathbb{R}^2 \times (-\infty, \infty)$ be a compact set. Without loss of generality we may assume that $K = \bar{B}_{r_0} \times [\tau_0, \tau'_0]$ for some $r_0 > 0$ and $\tau_0 < \tau'_0$. Then for any $|y| \leq r_0$ and $k \geq k_0$, we have

$$|y - \alpha_k^{-\frac{1}{2}} x_{t_k}| \leq r_0 + 2r_2.$$

Since

$$q_k(y, \tau) = \bar{u}_k(y - \alpha_k^{-\frac{1}{2}} x_{t_k}, \tau),$$

by Theorem 1.6,

$$q_k(y, \tau) - U(y - \alpha_k^{-\frac{1}{2}} x_{t_k}, \tau) \rightarrow 0 \quad \text{uniformly on } C^\infty(K) \quad \text{as } k \rightarrow \infty. \quad (1.37)$$

Now

$$\begin{aligned} |U(y, \tau) - U(y - \alpha_k^{-\frac{1}{2}} x_{t_k}, \tau)| &\leq \frac{\lambda}{(\lambda \rho_0 + e^{4\lambda\tau})^2} ||y - \alpha_k^{-\frac{1}{2}} x_{t_k}|^2 - |y|^2| \\ &\leq \lambda e^{-8\lambda\tau_0} \alpha_k^{-\frac{1}{2}} |x_{t_k}| (2|y| + \alpha_k^{-\frac{1}{2}} |x_{t_k}|) \\ &\leq 4\lambda e^{-8\lambda\tau_0} r_2 (r_0 + r_2) \alpha_k^{-\frac{1}{2}} \end{aligned}$$

holds for any $|y| \leq r_0$ and $\tau_0 \leq \tau \leq \tau'_0$ where ρ_0 is some constant between $|y|^2$ and $|y - \alpha_k^{-\frac{1}{2}} x_{t_k}|^2$. Similarly for any $\gamma_0, \gamma_1, \gamma_2 \in \mathbb{Z}^+ \cup \{0\}$,

$$|\partial_\tau^{\gamma_0} \partial_{y_1}^{\gamma_1} \partial_{y_2}^{\gamma_2} U(y, \tau) - \partial_\tau^{\gamma_0} \partial_{y_1}^{\gamma_1} \partial_{y_2}^{\gamma_2} U(y - \alpha_k^{-\frac{1}{2}} x_{t_k}, \tau)| \leq C \alpha_k^{-\frac{1}{2}} \quad (1.38)$$

holds for some constant $C > 0$ and any $|y| \leq r_0$, $\tau_0 \leq \tau \leq \tau'_0$. Since $\alpha_k \rightarrow \infty$ as $k \rightarrow \infty$ by Corollary 1.9, by (1.38)

$$U(y, \tau) - U(y - \alpha_k^{-\frac{1}{2}} x_{t_k}, \tau) \rightarrow 0 \quad \text{uniformly on } C^\infty(K) \quad \text{as } k \rightarrow \infty. \quad (1.39)$$

By (1.37) and (1.39) the theorem follows. $\square$

Theorem 1.11. Let $\{\tau_k\}_{k=1}^\infty$ be a sequence such that $\tau_k > 1/T$ for all $k \in \mathbb{Z}^+$ and $\tau_k \rightarrow \infty$ as $k \rightarrow \infty$. Let

$$\beta(\tau) = 1/\bar{u}(0, \tau), \quad (1.40)$$

$\beta_k = \beta(\tau_k)$ and

$$\bar{q}_k(y, \tau) = \beta_k \bar{u}(\beta_k^{\frac{1}{2}} y, \tau + \tau_k) \quad (1.41)$$

where $\bar{u}$ is given by (1.4). Then $\bar{q}_k$ has a subsequence $\bar{q}_{k_i}$ that converges uniformly on $C^\infty(K)$ for any compact set $K \subset \mathbb{R}^2 \times (-\infty, \infty)$ to some function $U(y, \tau)$ given by (1.23) for some $\lambda > 0$ as $\tau \rightarrow \infty$. Moreover $\beta_k \rightarrow \infty$ as $k \rightarrow \infty$.

Proof: Let $q_k(y, \tau)$ be given by (1.36) with α_k given by (1.11). Let $K \subset \mathbb{R}^2 \times (-\infty, \infty)$ be a compact set. As before we may assume without loss of generality that $K = \bar{B}_{r_0} \times [\tau_0, \tau'_0]$ for some $r_0 > 0$ and $\tau_0 < \tau'_0$. By Lemma 1.10 $q_k(y, \tau)$ has a subsequence which we may assume without loss of generality to be the sequence itself that converges uniformly on $C^\infty(K)$ as $k \rightarrow \infty$. Then

$$\frac{\alpha_k}{\beta_k} = q_k(0, 0) \rightarrow U(0, 0) = 1 \quad \text{as } k \rightarrow \infty. \quad (1.42)$$

Hence there exists $k_0 \in \mathbb{Z}^+$ and constants $c_2 > c_1 > 0$ such that

$$c_1 \leq \frac{\beta_k}{\alpha_k} \leq c_2 \quad \forall k \geq k_0. \quad (1.43)$$

Now

$$\bar{q}_k(y, \tau) = \frac{\beta_k}{\alpha_k} q_k((\beta_k/\alpha_k)^{\frac{1}{2}} y, \tau). \quad (1.44)$$

Then for any $|y| \leq r_0$, by (1.43),

$$(\beta_k/\alpha_k)^{\frac{1}{2}} |y| \leq c_2^{\frac{1}{2}} r_0 \quad \forall k \geq k_0.$$

Hence by Lemma 1.10

$$q_k((\beta_k/\alpha_k)^{\frac{1}{2}} y, \tau) \rightarrow U((\beta_k/\alpha_k)^{\frac{1}{2}} y, \tau) \quad (1.45)$$

uniformly on $C^\infty(K)$ as $k \rightarrow \infty$. By (1.23), (1.42), (1.44), (1.45) and Corollary 1.9 the theorem follows. $\square$

Lemma 1.12. Let τ_k , τ_{k_i} , β_k , $U(y, \tau)$ and $\lambda > 0$ be as in Theorem 1.11. Then for any $\varepsilon > 0$ and $M > 0$ there exist $n_1 \in \mathbb{Z}^+$ and $C > 0$ such that

$$\begin{cases} \left| u(x, t_{k_i}) - \frac{(T - t_{k_i})^2}{\lambda|x|^2 + \beta_{k_i}} \right| < u(0, t_{k_i})\varepsilon & \forall |x| \leq \beta_{k_i}^{\frac{1}{2}} M, i \geq n_1 \\ u(0, t_{k_i}) \leq C(T - t_{k_i})^2 & \forall i \geq n_1. \end{cases} \quad (1.46)$$

where t_k is given by (1.10).

Proof. Let $\bar{q}_k$ be given by (1.41) and k_0 be as in the proof of Theorem 1.11. By Theorem 1.11 $\bar{q}_k$ has a subsequence $\bar{q}_{k_i}$ that converges uniformly on $C^\infty(K)$ for any compact $K \subset \mathbb{R}^2 \times (-\infty, \infty)$ to some function $U(y, \tau)$ given by (1.23) as $k \rightarrow \infty$. Then there exists $n_1 \in \mathbb{Z}^+$ such that $k_i \geq k_0$ for all $i \geq n_1$ and

$$\begin{aligned} & \left| \bar{q}_{k_i}(y, 0) - \frac{1}{\lambda|y|^2 + 1} \right| < \varepsilon \quad \forall |y| \leq M, i \geq n_1 \\ \Rightarrow & \left| \beta_{k_i} \tau_{k_i}^2 u(x, t_{k_i}) - \frac{1}{\lambda \beta_{k_i}^{-1} |x|^2 + 1} \right| < \varepsilon \quad \forall |x| \leq \beta_{k_i}^{\frac{1}{2}} M, i \geq n_1 \\ \Rightarrow & \left| u(x, t_{k_i}) - \frac{(T - t_{k_i})^2}{\lambda |x|^2 + \beta_{k_i}} \right| < u(0, t_{k_i}) \varepsilon \quad \forall |x| \leq \beta_{k_i}^{\frac{1}{2}} M, i \geq n_1. \end{aligned}$$

By Corollary 1.8 and (1.43) there exists a constant $C > 0$ such that

$$\beta_{k_i} > C \quad \forall i \geq n_1$$

and the lemma follows. $\square$

Since the scalar curvature $R(x, t)$ satisfies

$$R_t = \Delta_{g(t)} R + R^2 \quad \text{in } \mathbb{R}^2 \times (0, T)$$

where $\Delta_{g(t)} = \frac{1}{u} \Delta$ is the Laplace-Beltrami operator with respect to the metric $g_{ij}(t)$, by (0.6) and the maximum principle,

$$R(x, t) \geq -\frac{1}{t + \mu} \quad \text{in } \mathbb{R}^2 \times (0, T). \quad (1.47)$$

Then by an argument similar to the proof of Lemma 3.5 of [DS2] but with Lemma 2.4, Lemma 3.4, and $k(t)$ in the proof there being replaced by Lemma 1.1, Lemma 1.12, and $1/[2(t + \mu)]$ we have the following lemma.

Lemma 1.13. *The constant λ in Theorem 1.11 satisfies*

$$\lambda \geq \frac{T + \mu}{2}.$$

By Lemma 1.13 and an argument similar to the proof of Lemma 3.5 of [DP2] we have the following lemma.

Lemma 1.14. *Let $\beta(\tau)$ be given by (1.40). Then*

$$\liminf_{\tau \rightarrow \infty} \frac{\beta'(\tau)}{\beta(\tau)} \geq 2(T + \mu).$$

Corollary 1.15. *Let $\beta(\tau)$ be given by (1.40). Then*

$$\beta(\tau) \geq e^{2(T+\mu)\tau + o(\tau)} \quad \text{as } \tau \rightarrow \infty.$$

As in [DS2] we consider the cylindrical change of variables,

$$v(\zeta, \theta, t) = r^2 u(r, \theta, t), \quad \zeta = \log r, r = |x|$$

and let

$$\tilde{v}(\xi, \theta, \tau) = \tau^2 v(\tau \xi, \theta, t), \quad \tau = 1/(T - t), \tau \geq 1/T. \quad (1.48)$$

Then $\tilde{v}$ satisfies

$$\tau \tilde{v}_\tau = \frac{1}{\tau} (\log \tilde{v})_{\xi\xi} + \tau (\log \tilde{v})_{\theta\theta} + \xi \tilde{v}_\xi + 2\tilde{v} \quad \text{in } \mathbb{R} \times [0, 2\pi] \times (1/T, \infty). \quad (1.49)$$

Lemma 1.16. *With the same notation as Theorem 1.11 for any $\varepsilon > 0$ there exists $n_1 \in \mathbb{Z}^+$ such that*

$$\left| \tilde{v}(\xi, \theta, \tau_{k_i}) - \frac{e^{2\tau_{k_i}\xi}}{\lambda e^{2\tau_{k_i}\xi} + \beta_{k_i}} \right| < \frac{e^{2\tau_{k_i}\xi}}{\beta_{k_i}} \varepsilon \quad \forall \xi \leq \frac{\log \beta_{k_i}}{2\tau_{k_i}}, \theta \in [0, 2\pi], i \geq n_1.$$

Proof: Let $\varepsilon > 0$. By Lemma 1.12 there exists $n_1 \in \mathbb{Z}^+$ such that (1.46) holds with $M = 1$. Since

$$\tilde{v}(\xi, \theta, \tau_{k_i}) = \tau_{k_i}^2 e^{2\tau_{k_i}\xi} u(e^{\tau_{k_i}\xi}, \theta, t_{k_i})$$

where t_{k_i} is given by (1.10) with $k = k_i$, by (1.46),

$$\left| \tilde{v}(\xi, \theta, \tau_{k_i}) - \frac{e^{2\tau_{k_i}\xi}}{\lambda e^{2\tau_{k_i}\xi} + \beta_{k_i}} \right| < \tau_{k_i}^2 e^{2\tau_{k_i}\xi} u(0, t_{k_i}) \varepsilon = \frac{e^{2\tau_{k_i}\xi}}{\beta_{k_i}} \varepsilon$$

holds for any $\xi \leq \log \beta_{k_i} / (2\tau_{k_i})$, $\theta \in [0, 2\pi]$ and $i \geq n_1$ and the lemma follows. $\square$

By (0.5), Corollary 1.15, Lemma 1.16, and an argument similar to the proof of Proposition 3.7 of [DP2] we have the following result.

Proposition 1.17. *Let $\beta(\tau)$ be given by (1.40). Then*

$$\lim_{\tau \rightarrow \infty} \frac{\log \beta(\tau)}{\tau} = 2(T + \mu).$$

Proposition 1.18. *Let $\beta(\tau)$ be given by (1.40). Then*

$$\lim_{\tau \rightarrow \infty} \frac{\beta'(\tau)}{\beta(\tau)} = 2(T + \mu). \quad (1.50)$$

Proof: Since

$$(\log \beta(\tau))_\tau = -(\log \bar{u}(0, \tau))_\tau = -\frac{\bar{u}_\tau(0, \tau)}{\bar{u}(0, \tau)}, \quad (1.51)$$

by (1.8) and (1.51),

$$C \geq \frac{\beta'(\tau)}{\beta(\tau)} \geq -\frac{2}{\tau^2 T} - \frac{2}{\tau} \geq -\frac{3T}{2} \quad \forall \tau \geq 2/T. \quad (1.52)$$

By (1.52) any sequence $\{\tau_k\}_{k=1}^\infty$, $\tau_k \rightarrow \infty$ as $k \rightarrow \infty$, will have a subsequence $\{\tau_{k_i}\}_{i=1}^\infty$ such that the limit

$$\lim_{i \rightarrow \infty} \frac{\beta'(\tau_{k_i})}{\beta(\tau_{k_i})}$$

exists. By the L'Hospital rule and Proposition 1.17,

$$\lim_{i \rightarrow \infty} \frac{\beta'(\tau_{k_i})}{\beta(\tau_{k_i})} = 2(T + \mu).$$

Since the sequence $\{\tau_k\}_{k=1}^\infty$ is arbitrary, (1.50) holds. $\square$

Proposition 1.19. *Let τ_k , $\bar{q}_k$, be as given in Theorem 1.11 and let*

$$\tilde{R}_k = -\frac{\Delta \log \bar{q}_k}{\bar{q}_k} \quad (1.53)$$

Then

$$\lim_{\tau \rightarrow \infty} \tilde{R}_k(0, 0) = 2(T + \mu).$$

Proof: By (1.5) $\bar{q}_k$ satisfies

$$\bar{q}_{k,\tau} = \Delta \log \bar{q}_k + \frac{2\bar{q}_k}{\tau + \tau_k} \quad \text{in } \mathbb{R}^2 \times (-\tau_k + (1/T), \infty). \quad (1.54)$$

By (1.41), (1.51) and (1.54),

$$\tilde{R}_k(0, 0) = \frac{\beta'(\tau_k)}{\beta(\tau_k)} + \frac{2}{\tau_k}. \quad (1.55)$$

Letting $k \rightarrow \infty$ in (1.55), by Proposition 1.18 the corollary follows. $\square$

Corollary 1.20. *The constant λ in Theorem 1.11 is equal to $(T + \mu)/2$.*

Proof: Let τ_k , $\bar{q}_k$, $\bar{q}_{k_i}$, and $U(y, \tau)$ be as in Theorem 1.11 and let $\tilde{R}_k$ be given by (1.53). Then by Theorem 1.11,

$$\lim_{i \rightarrow \infty} \tilde{R}_{k_i}(0, 0) = -\lim_{i \rightarrow \infty} \frac{\Delta \log \bar{q}_{k_i}}{\bar{q}_{k_i}}(0, 0) = -\frac{\Delta \log U}{U}(0, 0) = 4\lambda. \quad (1.56)$$

By Proposition 1.19 and (1.56) the corollary follows. $\square$

By Theorem 1.11 and Corollary 1.20 we have the following main theorem of this section.

Theorem 1.21. *Let $\beta(\tau)$ be given by (1.40) and let*

$$\bar{q}(y, \tau) = \beta(\tau) \bar{u}(\beta(\tau)^{\frac{1}{2}} y, \tau)$$

where $\bar{u}$ is given by (1.4). Then $\bar{q}(y, \tau)$ converges uniformly on $C^\infty(K)$ for any compact set $K \subset \mathbb{R}^2$ to the function

$$U_\mu(y) = \frac{1}{\frac{(T+\mu)}{2}|y|^2 + 1}$$

as $\tau \rightarrow \infty$.

Since

$$(\log \beta(\tau))_\tau = \bar{R}(0, \tau) - \frac{2}{\tau},$$

by (1.5), by Proposition 1.18 we have the following result.

Proposition 1.22. *Let $\beta(\tau)$ be given by (1.40). Then*

$$\lim_{\tau \rightarrow \infty} \bar{R}(0, \tau) = 2(T + \mu).$$

By Theorem 1.21 and an argument similar to the proof of Lemma 1.16 we have the following result.

Lemma 1.23. *Let $\beta(\tau)$ be given by (1.40) and let $\tilde{v}$ be given by (1.48). Then for any $\varepsilon > 0$ there exists $\tau_0 > 1/T$ such that*

$$\left| \tilde{v}(\xi, \theta, \tau) - \frac{e^{2\tau\xi}}{\frac{T+\mu}{2}e^{2\tau\xi} + \beta(\tau)} \right| < \frac{e^{2\tau\xi}}{\beta(\tau)} \varepsilon \quad \forall \xi \leq \frac{\log \beta(\tau)}{2\tau}, \theta \in [0, 2\pi], \tau \geq \tau_0.$$

By Corollary 1.15, Proposition 1.17 and Lemma 1.23 we get the following result.

Corollary 1.24. *Let $\beta(\tau)$ be given by (1.40) and let $\tilde{v}$ be given by (1.48). Then*

$$\int_{-\infty}^{\xi^-} \int_0^{2\pi} \tilde{v}(\xi, \theta, \tau) d\theta d\xi \rightarrow 0 \quad \text{as } \tau \rightarrow \infty$$

and

$$\lim_{\tau \rightarrow \infty} \tilde{v}(\xi, \theta, \tau) = 0 \quad \text{uniformly on } (-\infty, \xi^-] \times [0, 2\pi]$$

for any $\xi^- < T + \mu$.

2 Outer region behaviour

In this section we will prove the behaviour of the maximal solution in the outer region without using the Hamilton-Yau Harnack inequality for surfaces [HY]. By (0.5), Proposition 1.17, Lemma 1.23, Corollary 1.24 and an argument similar to the proof of Lemma 4.1 of [DS2] we have the following lemma.

Lemma 2.1. *Let $\xi(\tau) = (\log \beta(\tau))/2\tau$ with $\beta(\tau)$ given by (1.40). Let $\tilde{v}$ be given by (1.48). Then there exists constants $C_1 > 0$, $C_2 > 0$, $C_3 > 0$ and $\tau_0 > 1/T$ such that the following holds.*

- (i) $\tilde{v}(\xi, \theta, \tau) \leq C_1 \quad \forall \xi \in \mathbb{R}, \theta \in [0, 2\pi], \tau \geq 1/T$
- (ii) $\tilde{v}(\xi, \theta, \tau) \geq \frac{C_2}{\xi^2} \quad \forall \xi \geq \xi(\tau), \theta \in [0, 2\pi], \tau \geq \tau_0$
- (iii) $\tilde{v}(\xi, \theta, \tau) \leq \frac{C_3}{\xi^2} \quad \forall \xi > 0, \theta \in [0, 2\pi], \tau \geq \tau_0.$

Moreover

$$\xi(\tau) = T + \mu + o(1) \quad \text{as } \tau \rightarrow \infty.$$

Lemma 2.2. *For any $b > a > T + \mu$, there exist constants $C > 0$ and $\tau_1 > 1/T$ such that*

$$\max_{a \leq \xi \leq b} \left| \int_0^{2\pi} (\log \tilde{v})_\xi(\xi, \theta, \tau) d\theta \right| \leq C \quad \forall \tau \geq \tau_1. \quad (2.1)$$

Proof: Let $\delta = (a - (T + \mu))/2$. By direct computation the scalar curvature R in polar coordinates satisfies

$$\begin{aligned} R(e^{\tau\xi}, \theta, t) &= -\frac{(\log \tilde{v})_{\xi\xi} + \tau^2(\log \tilde{v})_{\theta\theta}}{\tilde{v}}(\xi, \theta, \tau) \\ \Rightarrow (\log \tilde{v})_{\xi\xi}(\xi, \theta, \tau) + \tau^2(\log \tilde{v})_{\theta\theta}(\xi, \theta, \tau) &= -R(e^{\tau\xi}, \theta, t)\tilde{v}(\xi, \theta, \tau) \end{aligned} \quad (2.2)$$

where $\tau = 1/(T - t)$. Integrating (2.2) over $(\theta, \xi) \in [0, 2\pi] \times [\xi_1, \xi_2]$, $T + \mu + \delta \leq \xi_1 < \xi_2 \leq b + 1$, by (1.47) and Lemma 2.1,

$$\begin{aligned} \int_0^{2\pi} (\log \tilde{v})_\xi(\xi_2, \theta, \tau) d\theta &= - \int_{\xi_1}^{\xi_2} \int_0^{2\pi} R(e^{\tau\xi}, \theta, t) \tilde{v}(\xi, \theta, \tau) d\theta d\xi \\ &\quad + \int_0^{2\pi} (\log \tilde{v})_\xi(\xi_1, \theta, \tau) d\theta \\ &\leq C_1 + \int_0^{2\pi} (\log \tilde{v})_\xi(\xi_1, \theta, \tau) d\theta \quad \forall \tau > 1/T \end{aligned} \quad (2.3)$$

for some constant $C_1 > 0$. Let $\tau_0 > 1/T$ be as given in Lemma 2.1. By Lemma 2.1 there exists $\tau_1 > \tau_0$ such that

$$\xi(\tau) < T + \mu + \delta \quad \forall \tau \geq \tau_1. \quad (2.4)$$

Integrating (2.3) over $\xi_1 \in (a - \delta, a)$, by (2.4) and Lemma 2.1,

$$\begin{aligned} \int_0^{2\pi} (\log \tilde{v})_\xi(\xi_2, \theta, \tau) d\theta &\leq \frac{1}{\delta} \left(C_1 + \int_0^{2\pi} (\log \tilde{v})(\xi_1, \theta, \tau) d\theta \Big|_{\xi_1=a-\delta}^{\xi_1=a} \right) \\ &\leq \frac{C'}{\delta} \quad \forall a \leq \xi_2 \leq b, \tau \geq \tau_1. \end{aligned} \quad (2.5)$$

Integrating (2.3) over $\xi_2 \in (b, b + 1)$, by (2.4) and Lemma 2.1,

$$\begin{aligned} \int_0^{2\pi} (\log \tilde{v})_\xi(\xi_1, \theta, \tau) d\theta &\geq -C_1 + \int_0^{2\pi} (\log \tilde{v})(\xi_1, \theta, \tau) d\theta \Big|_{\xi_1=b}^{\xi_1=b+1} \\ &\geq -C'' \quad \forall a \leq \xi_1 \leq b, \tau \geq \tau_1. \end{aligned} \quad (2.6)$$

By (2.5) and (2.6) we get (2.1) and the lemma follows. $\square$

We now let

$$w(\xi, \theta, s) = \tilde{v}(\xi, \theta, \tau) \quad (2.7)$$

with

$$s = \log \tau = -\log(T - t).$$

Then as in [DS2] by (1.49),

$$w_s = e^{-s}(\log w)_{\xi\xi} + e^s(\log w)_{\theta\theta} + \xi w_\xi + 2w \quad \text{in } \mathbb{R} \times [0, 2\pi] \times (-\log T, \infty). \quad (2.8)$$

Theorem 2.3. *As $\tau \rightarrow \infty$, the function $\tilde{v}$ given by (1.48) converges to the function*

$$V(\xi) = \begin{cases} \frac{2(T + \mu)}{\xi^2} & \forall \xi > T + \mu \\ 0 & \forall \xi < T + \mu. \end{cases}$$

Moreover the convergence is uniform on $(-\infty, a]$ for any $a < T + \mu$ and on $[\xi_0, \xi'_0]$ for any $\xi'_0 > \xi_0 > T + \mu$.

Proof: By Corollary 1.24 we only need to prove the convergence of the function $\tilde{v}$ to $2(T + \mu)/\xi^2$ for $\xi > T + \mu$. Let τ_0, τ_1 , be given by Lemma 2.1 and Lemma 2.2 respectively. Let $s_0 = \max(\log \tau_0, \log \tau_1)$ and $\{s_k\}_{k=1}^\infty$ be a sequence such that $s_k \rightarrow \infty$ as $k \rightarrow \infty$. Let

$$w_k(\xi, \theta, s) = w(\xi, \theta, s + s_k) \quad \forall \xi \in \mathbb{R}, 0 \leq \theta \leq 2\pi, s \geq -\log T - s_k. \quad (2.9)$$

Then by (2.8),

$$w_{k,s} = e^{-(s+s_k)}(\log w_k)_{\xi\xi} + e^{s+s_k}(\log w_k)_{\theta\theta} + \xi w_{k,\xi} + 2w_k \quad (2.10)$$

in $\mathbb{R} \times [0, 2\pi] \times (-\log T - s_k, \infty)$. Since

$$\int_{\mathbb{R}^2} u(x, t) dx = 4\pi(T - t) \quad \forall 0 < t < T,$$

by (1.48), (2.7) and (2.9),

$$\int_{-\infty}^{\infty} \int_0^{2\pi} w_k(\xi, \theta, s) d\theta d\xi = 4\pi \quad \forall s > -\log T - s_k, k \in \mathbb{Z}^+. \quad (2.11)$$

Let

$$W_k^b(\eta, s) = \int_{\eta}^b \int_0^{2\pi} w_k(\xi, \theta, s) d\theta d\xi \quad \forall b \geq \eta > T + \mu, s > -\log T - s_k, k \in \mathbb{Z}^+ \quad (2.12)$$

and

$$W_k(\eta, s) = \int_{\eta}^{\infty} \int_0^{2\pi} w_k(\xi, \theta, s) d\theta d\xi \quad \forall \eta > T + \mu, s > -\log T - s_k, k \in \mathbb{Z}^+.$$

Then

$$(W_k^b)_{\eta}(\eta, s) = - \int_0^{2\pi} w_k(\eta, \theta, s) d\theta \quad \forall b \geq \eta > T + \mu, s > -\log T - s_k, k \in \mathbb{Z}^+. \quad (2.13)$$

Hence by (2.11) and Lemma 2.1 there exists a constant $C > 0$ such that

$$|W_k^b(\eta, s)|, |(W_k^b)_{\eta}(\eta, s)| \leq C \quad \forall b \geq \eta > T + \mu, s > s_0 - s_k, k \in \mathbb{Z}^+. \quad (2.14)$$

Now by (2.10), (2.12) and (2.13),

$$\begin{aligned} (W_k^b)_s(\eta, s) &= e^{-(s+s_k)} \int_{\eta}^b \int_0^{2\pi} (\log w_k)_{\xi\xi}(\xi, \theta, s) d\theta d\xi + \int_{\eta}^b \int_0^{2\pi} \xi w_{k,\xi}(\xi, \theta, s) d\theta d\xi \\ &\quad + 2W_k^b \\ &= e^{-(s+s_k)} \int_0^{2\pi} (\log w_k)_{\xi}(b, \theta, s) d\theta - e^{-(s+s_k)} \int_0^{2\pi} (\log w_k)_{\xi}(\eta, \theta, s) d\theta \\ &\quad + b \int_0^{2\pi} w_k(b, \theta, s) d\theta - \eta \int_0^{2\pi} w_k(\eta, \theta, s) d\theta + W_k^b \\ &= e^{-(s+s_k)} \int_0^{2\pi} (\log w_k)_{\xi}(b, \theta, s) d\theta - e^{-(s+s_k)} \int_0^{2\pi} (\log w_k)_{\xi}(\eta, \theta, s) d\theta \\ &\quad + b \int_0^{2\pi} w_k(b, \theta, s) d\theta + \eta(W_k^b)_{\eta}(\eta, s) + W_k^b. \end{aligned} \quad (2.15)$$

By (2.14), Lemma 2.1, and Lemma 2.2 for any $a \in (T + \mu, b]$ there exists a constant $C_{a,b} > 0$ such that

$$|(W_k^b)_s(\eta, s)| \leq C_{a,b}(1 + b^{-1} + \eta) \quad \forall a \leq \eta \leq b, s > s_0 - s_k, k \in \mathbb{Z}^+. \quad (2.16)$$

By (2.14) and (2.16) the sequence $\{W_k^b\}_{k=1}^\infty$ is equi-Holder continuous on $[a, b] \times [-s', \infty]$ for any $a \in (T + \mu, b]$, $s' > 0$, and k large such that $s_k - s_0 > s'$.

We choose a sequence $\{b_i\}_{i=1}^\infty$ of monotonically increasing sequence such that $b_i > T + \mu$ for any $i \in \mathbb{Z}^+$ and $b_i \rightarrow \infty$ as $i \rightarrow \infty$. By the Ascoli theorem and a diagonalization argument the sequence $\{W_k^{b_1}\}_{k=1}^\infty$ has a subsequence $\{W_{j_{1,k}}^{b_1}\}_{k=1}^\infty$ such that $W_{j_{1,k}}^{b_1}$ converges uniformly on every compact subset of $(T + \mu, b_1] \times (-\infty, \infty)$ to some function $W^{b_1} \in C((T + \mu, b_1] \times (-\infty, \infty))$ as $k \rightarrow \infty$.

Similarly the sequence $\{W_{j_{1,k}}^{b_2}\}_{k=1}^\infty$ has a subsequence $\{W_{j_{2,k}}^{b_2}\}_{k=1}^\infty$ with $j_{1,1} < j_{2,1}$ such that $W_{j_{2,k}}^{b_2}$ converges uniformly on every compact subset of $(T + \mu, b_2] \times (-\infty, \infty)$ to some function $W^{b_2} \in C((T + \mu, b_2] \times (-\infty, \infty))$ as $k \rightarrow \infty$. Repeating the above argument for any $i \geq 2$ the sequence $\{W_{j_{i-1,k}}^{b_i}\}_{k=1}^\infty$ has a subsequence $\{W_{j_{i,k}}^{b_i}\}_{k=1}^\infty$ with $j_{i-1,1} < j_{i,1}$ such that $W_{j_{i,k}}^{b_i}$ converges uniformly on every compact subset of $(T + \mu, b_i] \times (-\infty, \infty)$ to some function $W^{b_i} \in C((T + \mu, b_i] \times (-\infty, \infty))$ as $k \rightarrow \infty$.

For any $k \in \mathbb{Z}^+$ let $j_k = j_{k,1}$. Then $\{j_k\}_{k=1}^\infty$ is a subsequence of $\{j_{i,k}\}_{k=1}^\infty$ for any $i \in \mathbb{Z}^+$. Hence for any $i \in \mathbb{Z}^+$, the sequence $\{W_{j_k}^{b_i}\}_{k=1}^\infty$ converges uniformly on every compact subset of $(T + \mu, b_i] \times (-\infty, \infty)$ to W^{b_i} as $k \rightarrow \infty$. Thus we may assume without loss of generality that for any $i \in \mathbb{Z}^+$ $\{W_k^{b_i}\}_{k=1}^\infty$ converges uniformly on every compact subset of $(T + \mu, b_i] \times (-\infty, \infty)$ to W^{b_i} as $k \rightarrow \infty$.

Now by Lemma 2.1 for each $s \in \mathbb{R}$, $w_k(\cdot, \cdot, s)$ will have a subsequence which we may assume without loss of generality to be the sequence itself that converges weakly in $L^\infty(K)$ to some non-negative function $\bar{w}(\cdot, \cdot, s)$ as $k \rightarrow \infty$ for any compact set $K \subset [T + \mu, \infty) \times [0, 2\pi]$. Putting $b = b_i$ and letting $k \rightarrow \infty$ in (2.12),

$$W^{b_i}(\eta, s) = \int_\eta^{b_i} \int_0^{2\pi} \bar{w}(\xi, \theta, s) d\theta d\xi \quad \forall b \geq \eta \geq T + \mu, s \in \mathbb{R}, i \in \mathbb{Z}^+. \quad (2.17)$$

By (2.17) as $i \rightarrow \infty$, W^{b_i} will increase monotonically to the function,

$$W(\eta, s) := \int_\eta^\infty \int_0^{2\pi} \bar{w}(\xi, \theta, s) d\theta d\xi \quad \forall \eta \geq T + \mu, s \in \mathbb{R}. \quad (2.18)$$

By Lemma 2.1,

$$\bar{w}(\xi, \theta, s) \leq \frac{C}{\xi^2} \quad \forall \xi > 0, \theta \in [0, 2\pi], s \in \mathbb{R}. \quad (2.19)$$

Then by (2.19),

$$0 \leq W(\eta, s) - W^{b_i}(\eta, s) \leq \int_{b_i}^\infty \int_0^{2\pi} \bar{w}(\xi, \theta, s) d\theta d\xi \leq \int_{b_i}^\infty \int_0^{2\pi} \frac{C}{\xi^2} d\theta d\xi \leq \frac{C'}{b_i} \quad (2.20)$$

holds for any $s \in \mathbb{R}$ and $i \in \mathbb{Z}^+$.

Claim 1: W_k converges uniformly to W on every compact set $K \subset (T + \mu, \infty) \times (-\infty, \infty)$ as $k \rightarrow \infty$.

Proof of Claim 1: Let K be a compact subset of $(T + \mu, \infty) \times (-\infty, \infty)$. By Lemma 2.1, (2.19) and (2.20),

$$\begin{aligned} \|W_k - W\|_{L^\infty(K)} &\leq \|W_k - W_k^{b_i}\|_{L^\infty(K)} + \|W_k^{b_i} - W^{b_i}\|_{L^\infty(K)} + \|W^{b_i} - W\|_{L^\infty(K)} \\ &\leq \int_{b_i}^\infty \int_0^{2\pi} \frac{C}{\xi^2} d\theta d\xi + \|W_k^{b_i} - W^{b_i}\|_{L^\infty(K)} + \frac{C'}{b_i} \\ &\leq \frac{C''}{b_i} + \|W_k^{b_i} - W^{b_i}\|_{L^\infty(K)} \quad \forall i \in \mathbb{Z}^+. \end{aligned} \quad (2.21)$$

Letting $k \rightarrow \infty$ in (2.21),

$$\begin{aligned} \limsup_{k \rightarrow \infty} \|W_k - W\|_{L^\infty(K)} &\leq \frac{C''}{b_i} \quad \forall i \in \mathbb{Z}^+ \\ \Rightarrow \lim_{k \rightarrow \infty} \|W_k - W\|_{L^\infty(K)} &= 0 \quad \text{as } i \rightarrow \infty \end{aligned}$$

and Claim 1 follows.

By Corollary 1.24, Lemma 2.1, (2.7), (2.9), (2.11) and Claim 1,

$$W((T + \mu)^+, s) = 4\pi \quad \forall s \in \mathbb{R}. \quad (2.22)$$

Let $a_2 > a_1 > T + \mu$ and $s'_1 < s'_2$. We choose $k_0 \in \mathbb{Z}^+$ such that $s'_1 > s_0 - s_k$ for any $k \geq k_0$. Then by (2.15) for any $e^{-s}\zeta_0 \in [a_1, a_2]$, $b > a_2$, $s'_2 \geq s \geq s'_1$ and $k \geq k_0$,

$$\begin{aligned} &\frac{d}{ds}(e^{-s}W_k^b(e^{-s}\zeta_0, s)) \\ &= e^{-s}(W_{k,s}^b(e^{-s}\zeta_0, s) - e^{-s}\zeta_0 W_{k,\eta}^b(e^{-s}\zeta_0, s) - W_k^b(e^{-s}\zeta_0, s)) \\ &= e^{-(2s+s_k)} \int_0^{2\pi} (\log w_k)_\xi(b, \theta, s) d\theta - e^{-(2s+s_k)} \int_0^{2\pi} (\log w_k)_\xi(e^{-s}\zeta_0, \theta, s) d\theta \\ &\quad + be^{-s} \int_0^{2\pi} w_k(b, \theta, s) d\theta \end{aligned} \quad (2.23)$$

By Lemma 2.1 and Lemma 2.2,

$$\left| \frac{d}{ds}(e^{-s}W_k^b(e^{-s}\zeta_0, s)) \right| \leq C_b e^{-(2s+s_k)} + C \frac{e^{-s}}{b} \quad (2.24)$$

for some constants $C_b > 0$ depending on b and $C > 0$. By (2.24) for any $s'_2 > s'_1 > s_0 - s_k$, $\zeta_0 > e^{s'_2}(T + \mu)$ and $b > e^{-s'_1}\zeta_0$,

$$\begin{aligned} |e^{-s'_1}W_k^b(e^{-s'_1}\zeta_0, s'_1) - e^{-s'_2}W_k^b(e^{-s'_2}\zeta_0, s'_2)| &= \left| \int_{s'_1}^{s'_2} \frac{d}{ds}(e^{-s}W_k^b(e^{-s}\zeta_0, s)) ds \right| \\ &\leq \max_{s'_1 \leq s \leq s'_2} \left| \frac{d}{ds}(e^{-s}W_k^b(e^{-s}\zeta_0, s)) \right| (s'_2 - s'_1) \\ &\leq \left(C_b e^{-(2s'_1+s_k)} + C \frac{e^{-s'_1}}{b} \right) (s'_2 - s'_1) \end{aligned} \quad (2.25)$$

Putting $b = b_i$ and letting first $k \rightarrow \infty$ and then $i \rightarrow \infty$ in (2.25),

$$e^{-s'_1} W(e^{-s'_1} \zeta_0, s'_1) = e^{-s'_2} W(e^{-s'_2} \zeta_0, s'_2) \quad (2.26)$$

holds for any $s'_1, s'_2 \in \mathbb{R}$ and $\zeta_0 > \max(e^{s'_1}, e^{s'_2})(T + \mu)$.

Let $\eta, \bar{\eta} > T + \mu$ and $s \in \mathbb{R}$. Let $\zeta_0 = e^s \eta$ and choose $\bar{s}$ such that $\bar{\eta} = e^{-\bar{s}} \zeta_0$. Then $\eta/\bar{\eta} = e^{\bar{s}-s}$. Hence by (2.26),

$$\eta W(\eta, s) = \bar{\eta} W(\bar{\eta}, \bar{s}) \quad \forall \eta, \bar{\eta} > T + \mu, s, \bar{s} \in \mathbb{R}. \quad (2.27)$$

Letting $\bar{\eta} \rightarrow T + \mu$ in (2.27), by (2.22) we get

$$W(\eta, s) = \frac{4\pi(T + \mu)}{\eta} \quad \forall \eta > T + \mu, s \in \mathbb{R}. \quad (2.28)$$

We now fix $s \in \mathbb{R}$. By (2.28) and an argument similar to the proof on P.588 of [DS2] $w_k(\xi, \theta, s)$ converges to $2(T + \mu)/\xi^2$ in $L^p([\xi_0, \infty) \times [0, 2\pi])$ as $k \rightarrow \infty$ for any $p \geq 1$ and $\xi_0 > T + \mu$. Hence by passing to a subsequence we may assume without loss of generality that

$$w_k(\xi, \theta, s) \rightarrow \frac{2(T + \mu)}{\xi^2} \quad \text{a.e. } (\xi, \theta) \in (T + \mu, \infty) \times [0, 2\pi] \quad \text{as } k \rightarrow \infty. \quad (2.29)$$

Let

$$Z_k(\xi, s) = \int_0^{2\pi} \log w_k(\xi, \theta, s) d\theta.$$

Claim 2: For any $s \in \mathbb{R}$ the sequence $Z_k(\xi, s)$ has a subsequence that converges uniformly on $[a, b]$ for any $b > a > T + \mu$ to

$$Z(\xi, s) = 2\pi \log \frac{2(T + \mu)}{\xi^2}. \quad (2.30)$$

Proof of Claim 2: We will use a modification of the proof of Lemma 4.10 of [DS2] to prove the claim. By Lemma 2.1 and Lemma 2.2 for any $b > a > T + \mu$ there exist constants $C > 0$ and $k_1 = k_1(s) \in \mathbb{Z}^+$ such that

$$\begin{cases} |Z_{k,\xi}(\xi, s)| = \left| \int_0^{2\pi} (\log w_k(\xi, \theta, s))_\xi d\theta \right| \leq C & \forall a \leq \xi \leq b, k \geq k_1 \\ |Z_k(\xi, s)| \leq C & \forall a \leq \xi \leq b, k \geq k_1. \end{cases} \quad (2.31)$$

By (2.31), the Ascoli theorem, and a diagonalization argument the sequence $Z_k(\xi, \theta, s)$ has a subsequence that converges uniformly on $[a, b]$ for any $b > a > T + \mu$ to some function $Z(\xi, s)$ which is continuous in $\xi > T + \mu$. Without loss of generality we may

assume that $Z_k(\xi, s)$ converges uniformly on $[a, b]$ for any $b > a > T + \mu$ to $Z(\xi, s)$ as $k \rightarrow \infty$. Since

$$\int_{\eta}^{\eta+A} \int_0^{2\pi} \log w_k(\xi, \theta, s) d\theta d\xi \rightarrow \int_{\eta}^{\eta+A} Z(\xi, s) d\xi \quad \forall \eta > T + \mu, A > 0 \quad \text{as } k \rightarrow \infty,$$

by (2.29), Lemma 2.1, and the Lebesgue dominated convergence theorem,

$$\int_{\eta}^{\eta+A} Z(\xi, s) d\xi = \int_{\eta}^{\eta+A} \int_0^{2\pi} \log \frac{2(T + \mu)}{\xi^2} d\theta d\xi = 2\pi \int_{\eta}^{\eta+A} \log \frac{2(T + \mu)}{\xi^2} d\xi \quad (2.32)$$

holds for any $\eta > T + \mu$, $A > 0$. Dividing both sides of (2.32) by A and letting $A \rightarrow 0$ we get (2.30) and Claim 2 follows.

By Lemma 1.1,

$$\begin{aligned} w_k(\eta, \theta, s) &= \tilde{v}(\eta, \theta, \tau_k(s)) = \tau_k(s)^2 v(\tau_k(s)\eta, \theta, \tau_k(s)) \\ &= \tau_k(s)^2 e^{2\tau_k(s)\eta} u(e^{\tau_k(s)\eta}, \theta, t_k(s)) \\ &\leq \tau_k(s)^2 e^{2\tau_k(s)\eta} u(e^{\tau_k(s)\eta} - 2r_2, \theta, t_k(s)) \\ &= \tau_k(s)^2 e^{2\tau_k(s)\eta} u(e^{\tau_k(s)\eta}(1 - 2r_2 e^{-\tau_k(s)\eta}), \theta, t_k(s)) \\ &= \frac{\tau_k(s)^2}{(1 - 2r_2 e^{-\tau_k(s)\eta})^2} v(\tau_k(s)\eta + \log(1 - 2r_2 e^{-\tau_k(s)\eta}), \theta, t_k(s)) \\ &= (1 - 2r_2 e^{-\tau_k(s)\eta})^{-2} \tilde{v}(\eta + \tau_k(s)^{-1} \log(1 - 2r_2 e^{-\tau_k(s)\eta}), \theta, \tau_k(s)) \\ &= (1 - 2r_2 e^{-\tau_k(s)\eta})^{-2} w_k(\eta + \tau_k(s)^{-1} \log(1 - 2r_2 e^{-\tau_k(s)\eta}), \theta, s) \end{aligned} \quad (2.33)$$

for all k large satisfying $e^{\tau_k(s)\eta} > 2r_2$ where

$$\tau_k(s) = e^{s+s_k} \quad \text{and } t_k(s) = T - \tau_k(s)^{-1}.$$

By Claim 2, (2.33), and an argument similar to the proof on P.590 of [DS2], $w_k(\eta, \theta, s)$ converges uniformly on every compact subset of $(T + \mu, \infty) \times [0, 2\pi]$ to

$$\frac{2(T + \mu)}{\eta^2}$$

as $k \rightarrow \infty$. Since the sequence s_k is arbitrary, $\tilde{v}(\eta, \theta, \tau)$ converges uniformly to $2(T + \mu)/\eta^2$ on every compact subset of $(T + \mu, \infty) \times [0, 2\pi]$ as $\tau \rightarrow \infty$ and the theorem follows. $\square$

References

[CF] D.G. Aronson and L.A. Caffarelli, *The initial trace of a solution of the porous medium equation*, Transactions A.M.S. 280 (1983), no. 1, 351–366.

- [DH] P. Daskalopoulos and R. Hamilton, *Geometric estimates for the logarithmic fast diffusion equation*, Comm. Anal. Geom. 12 (2004), nos. 1–2, 143–164.
- [DP1] P. Daskalopoulos and M.A. del Pino, *On a singular diffusion equation*, Comm. Anal. Geom. 3 (1995), no. 3, 523–542.
- [DP2] P. Daskalopoulos and M.A. del Pino, *Type II collapsing of maximal solutions to the Ricci flow in $\mathbb{R}^2$* , Ann. Inst. H. Poincaré Anal. Non Linéaire 24 (2007), 851–874.
- [DS1] P. Daskalopoulos and N. Sesum, *Eternal solutions to the Ricci flow on $\mathbb{R}^2$* , Int. Math. Res. Not. 2006, Art. ID 83610, 20 pp.
- [DS2] P. Daskalopoulos and N. Sesum, *Type II extinction profile of maximal solutions to the Ricci flow equation*, J. Geom. Anal. 20 (2010), no. 3, 565–591.
- [ERV1] J.R. Esteban, A. Rodriguez and J.L. Vazquez, *The fast diffusion equation with logarithmic nonlinearity and the evolution of conformal metrics in the plane*, Advances in Differential Equations 1 (1996), no. 1, 21–50.
- [ERV2] J.R. Esteban, A. Rodriguez and J.L. Vazquez, *The maximal solution of the logarithmic fast diffusion equation in two space dimensions*, Advances in Differential Equations 2 (1997), no. 6, 867–894.
- [G] P.G. de Gennes, *Wetting: statics and dynamics*, Rev. Modern Phys. 57 (1985), no. 3, 827–863.
- [HY] R. Hamilton and S.T. Yau, *The Harnack estimate for the Ricci flow - revisited*, Asian J. Math 1 (1997), no. 3, 418–421.
- [Hs1] S.Y. Hsu, *Large time behaviour of solutions of the Ricci flow equation on R^2* , Pacific J. Math 197 (2001), no. 1, 25–41.
- [Hs2] S.Y. Hsu, *Asymptotic profile of a singular diffusion equation as $t \rightarrow \infty$* , Nonlinear Analysis TMA 48 (2002), no. 6, 781–790.
- [Hs3] S.Y. Hsu, *Asymptotic behaviour of solutions of the equation $u_t = \Delta \log u$ near the extinction time*, Advances in Differential Equations 8 (2003), no. 2, 161–187.
- [Hs4] S.Y. Hsu, *Behaviour of solutions of a singular diffusion equation near the extinction time*, Nonlinear Analysis TMA 56 (2004), no. 1, 63–104.
- [Hu1] K.M. Hui, *Existence of solutions of the equation $u_t = \Delta \log u$* , Nonlinear Analysis, TMA 37 (1999), no. 7, 875–914.
- [Hu2] K.M. Hui, *Singular limit of solutions of the equation $u_t = \Delta(u^m/m)$ as $m \rightarrow 0$* , Pacific J. Math. 187 (1999), no. 2, 297–316.

- [K] J.R. King, *Self-similar behaviour for the equation of fast nonlinear diffusion*, Phil. Trans. Royal Soc. London, Series A 343 (1993), 337–375.
- [LSU] O.A. Ladyzenskaya, V.A. Solonnikov, and N.N. Uraltceva, *Linear and quasi-linear equations of parabolic type*, Transl. Math. Mono. Vol 23, Amer. Math. Soc., Providence, R.I., U.S.A., 1968.
- [V] J.L. Vazquez, *Nonexistence of solutions for nonlinear heat equations of fast-diffusion type*, J. Math. Pures Appl. 71 (1992), 503–526.
- [W1] L.F. Wu, *A new result for the porous medium equation derived from the Ricci flow*, Bull. Amer. Math. Soc. 28 (1993), 90–94.
- [W2] L.F. Wu, *The Ricci flow on R^2* , Comm. Anal. Geom. 1 (1993), 439–472.